

9

Non-radioactive Microsphere Bibliography

Abe, Y., Y. Kitada and A. Narimatsu. Effect of a calcium-sensitizing positive inotropic agent MCI-154 and its combined use with enalapril on postischemic contractile dysfunction of dog hearts. *J Cardiovasc Pharmacol.* 26:653-9., 1995.

Abel, F. L., R. R. Beck and R. H. Cooper. Fluorescent microspheres [letter]. *J Appl Physiol.* 76:986, 1994.

Abel, F. L., R. H. Cooper and R. R. Beck. Use of fluorescent latex microspheres to measure coronary blood flow distribution. *Circ Shock.* 41:156-61, 1993.

Abraham, S. A., F. N. Mirecki, D. Levine, A. D. Nunn, H. W. Strauss and H. Gewirtz. Myocardial technetium-99m-teboroxime activity in acute coronary artery occlusion and reperfusion: relation to myocardial blood flow and viability. *J Nucl Med.* 36:1062-8., 1995.

Altemeier, W. A., H. T. Robertson and R. W. Glenny. Pulmonary gas-exchange analysis by using simultaneous deposition of aerosolized and injected microspheres. *Journal of Applied Physiology.* 85:2344-2351, 1998.

Altemeier, W. A., H. T. Robertson, S. McKinney and R. W. Glenny. Pulmonary embolization causes hypoxemia by redistributing regional blood flow without changing ventilation. *Journal of Applied Physiology.* 85:2337-2343, 1998.

Angelborg, C., N. Slepcky, H. C. Larsen and L. Soederberg. Colored microspheres for blood flow determinations twice in the same animal. *Heart Res.* 27:265-9, 1987.

Ardehali, A., R. N. Gates, H. Laks, D. C. Drinkwater, Jr., E. Rudis, T. J. Sorensen, P. Chang and A. Aharon. The regional capillary distribution of retrograde blood cardioplegia in explanted human hearts. *J Thorac Cardiovasc Surg.* 109:935-9; discussion 939-40., 1995.

Ardevol, A., C. Adan, X. Remesar, M. Alemany and J. A. Fernandezlopez. Muscle blood flow during intense exercise in the obese rat. *Archives of Physiology & Biochemistry.* 104:337-343, 1996.

Ardevol, A., C. Adan, X. Remesar, J.-A. FernandezLopez and M. Alemany. Hind leg heat balance in obese Zucker rats during exercise. *PFLUGERS ARCHIV-EUROPEAN JOURNAL OF PHYSIOLOGY.* 435:454-464, 1998.

Ardevol, A., C. A. X. Remesar, J. A. FernandezLopez and M. Alemany. Lactate-bicarbonate interrelationship during exercise and recovery in lean and obese Zucker rats. *International Journal Of Obesity.* 21:333-339, 1997.

Arshady, R. Microspheres for biomedical applications: preparation of reactive and labelled microspheres. *Biomaterials*. 14:5-15, 1993.

Atef, N., A. Ktorza, L. Picon and L. P'Enicaud. Increased islet blood flow in obese rats: role of the autonomic nervous system. *Am J Physiol*. 262:E736-40., 1992.

Atef, N., M. C. Laury, J. M. Nguyen, N. Mokhtar, A. Ktorza and L. Penicaud. Increased pancreatic islet blood flow in 48-hour glucose-infused rats: Involvement of central and autonomic nervous systems. *Endocrinology*. 138:1836-1840, 1997.

Atef, N., B. Portha and L. P'Enicaud. Changes in islet blood flow in rats with NIDDM. *Diabetologia*. 37:677-80., 1994.

Audet, D. M. and W. L. Olbricht. The motion of model cells at capillary bifurcations. *Microvasc Res*. 33:377-96, 1987.

Austin, G. E., M. B. Tuvlin, D. Martino-Salzman, R. L. Hunter, A. G. Justicz, N. K. Thompson and A. C. Brooks. Determination of regional myocardial blood flow using fluorescent microspheres. *Am J Cardiovasc Pathol*. 4:352-7, 1993.

Austin, R. E., Jr., W. W. Hauck, G. S. Aldea, A. E. Flynn, D. L. Coggins and J. I. Hoffman. Quantitating error in blood flow measurements with radioactive microspheres. *Am J Physiol*. 257:H280-8, 1989.

Baer, R. W., B. D. Payne, E. D. Verrier, G. J. Vlahakes, D. Molodowitch, P. N. Uhlig and J. I. E. Hoffman. Increased number of myocardial blood flow measurements with radionuclide-labeled microspheres. *Am. J. Physiol*. 246:H418-H434, 1994.

Baile, E. M., J. M. Nelems, M. Schulzer and P. D. Par'e. Measurement of regional bronchial arterial blood flow and bronchovascular resistance in dogs. *J Appl Physiol*. 53:1044-9, 1982.

Bailey, W. F., Jr., M. G. Magno, P. D. Buckman, F. DiMeo, T. Langan, V. T. Armenti and J. D. Mannion. Chronic stimulation enhances extramyocardial collateral blood flow after a cardiomyoplasty. *Ann Thorac Surg*. 56:1045-52; discussion 1052-3., 1993.

Barnes, M. D., N. Lermer, W. B. Whitten and J. M. Ramsey. CCD based approach to high-precision size and refractive index determination of levitated microdroplets using Fraunhofer diffraction. *Review Of Scientific Instruments*. 68:2287-2291, 1997.

Bassingthwaigte, J. B., M. A. Malone, T. C. Moffett, R. B. King, I. S. Chan, J. M. Link and K. A. Krohn. Molecular and particulate depositions for regional myocardial flows in sheep. *Circ Res*. 66:1328-44, 1990.

Bassingthwaigte, J. B., M. A. Malone, T. C. Moffett, R. B. King, S. E. Little, J. M. Link and K. A. Krohn. Validity of microsphere deposition for regional myocardial flows. *Am. J. Physiol*. 253:H184-H193, 1987.

Bauer, R., R. Bergmann, B. Walter, P. Brust, U. Zwiener and B. Johannsen. Regional distribution of cerebral blood volume and cerebral blood flow in newborn piglets - effect of hypoxia/hypercapnia. *Developmental Brain Research.* 112:89-98, 1999.

Bauer, R., D. Hoyer, B. Walter, E. Gaser, H. Kluge and U. Zwiener. Changed systemic and cerebral hemodynamics and oxygen supply due to gradual hemorrhagic hypotension induced by an external PID-controller in newborn swine. *EXPERIMENTAL AND TOXICOLOGIC PATHOLOGY.* 49:469-476, 1997.

Bauer, R., B. Walter, A. Torossian, H. Fritz, O. Schlonski, T. Jochum, D. Hoyer, K. Reinhart and U. Zwiener. A piglet model for evaluation of cerebral blood flow and brain oxidative metabolism during gradual cerebral perfusion pressure decrease. *Pediatric Neurosurgery.* 30:62-69, 1999.

Bauer, R., B. Walter, E. Wurker, H. Kluge and U. Zwiener. Colored microsphere technique as a new method for quantitative-multiple estimation of regional hepatic and portal blood flow. *Experimental & Toxicologic Pathology.* 48:415-420, 1996.

Baumgart, D., T. Ehring and G. Heusch. A proischaemic action of nisoldipine: relationship to a decrease in perfusion pressure and comparison to dipyridamole. *Cardiovasc Res.* 27:1254-9, 1993.

Baumgart, D., T. Ehring, P. Kowallik, B. D. Guth, M. Krajcar and G. Heusch. Impact of alpha-adrenergic coronary vasoconstriction on the transmural myocardial blood flow distribution during humoral and neuronal adrenergic activation. *Circ Res.* 73:869-86., 1993.

Beck, K. C. Regional trapping of microspheres in the lung compares well with regional blood flow. *J Appl Physiol.* 63:883-9, 1987.

Beezhold, D. H., G. L. Sussman, G. M. Liss and N. S. Chang. Latex Allergy Can Induce Clinical Reactions to Specific Foods. *Clinical & Experimental Allergy.* 26:416-422, 1996.

Bernard, S. L., R. W. Glenny, H. H. Erickson, M. R. Fedde, N. Polissar, R. J. Basaraba and M. P. Hlastala. Minimal redistribution of pulmonary blood flow with exercise in racehorses. *Journal Of Applied Physiology.* 81:1062-1070, 1996.

Bernard, S. L., R. W. Glenny, N. L. Polissar, D. L. Luchtel and S. Lakshminarayan. Distribution of pulmonary and bronchial blood supply to airways measured by fluorescent microspheres. *Journal of Applied Physiology.* 80:430-436, 1996.

Bernard, S. L., D. L. Luchtel, R. W. Glenny and S. Lakshminarayan. Bronchial circulation in the marsupial opossum, *Didelphis marsupialis*. *Respiration Physiology.* 105:1-2, 1996.

Black, S. C. and I. W. Rodger. Methods for studying experimental myocardial ischemic and reperfusion injury. *Journal Of Pharmacological And Toxicological Methods.* 35:179-190, 1996.

Bland, J. M. and D. G. Altman. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet.* 1:307-10, 1986.

Bock, J. C., A. H. G. Goetze and R. Felix. Confirmation by animal experiment of a semiquantitative single-layer tomography for MR imaging of regional cerebral perfusion. *Rofo Fortschritte Auf Dem Gebiet Der Rontgenstrahlen Und Der Bildgebenden Verfahren.* 166:335-341, 1997.

Boeckxstaens, C. J. and W. J. Flameng. Retrograde cerebral perfusion does not perfuse the brain in nonhuman primates [see comments]. *Ann Thorac Surg.* 60:319-27; discussion 327-8., 1995.

Bolender, R. P., D. M. Hyde and R. T. Dehoff. Lung morphometry: a new generation of tools and experiments for organ, tissue, cell, and molecular biology. *Am J Physiol.* 265:L521-48, 1993.

Bone, H. G., P. J. Schenarts, S. R. Fischer, R. McGuire, L. D. Traber and D. L. Traber. Pyridoxalated hemoglobin polyoxyethylene conjugate reverses hyperdynamic circulation in septic sheep. *JOURNAL OF APPLIED PHYSIOLOGY.* 84:1991-1999, 1998.

Booke, M., F. Hinder, R. McGuire, L. D. Traber and D. L. Traber. Selective inhibition of inducible nitric oxide synthase: Effects on hemodynamics and regional blood flow in healthy and septic sheep. *Critical Care Medicine.* 27:162-167, 1999.

Bouaziz, H., N. Okubo, J. M. Malinovsky, D. Benhamou, K. Samii and J. X. Mazoit. The age-related effects of epidural lidocaine, with and without epinephrine, on spinal cord blood flow in anesthetized rabbits. *Anesthesia and Analgesia.* 88:1302-1307, 1999.

Boyde, A., L. A. Wolfe, M. Maly and S. J. Jones. Vital confocal microscopy in bone. *Scanning.* 17:72-85, 1995.

Bray, R., K. Forrester, J. J. McDougall, A. Damji and W. R. Ferrell. Evaluation of laser Doppler imaging to measure blood flow in knee ligaments of adult rabbits. *Medical & Biological Engineering & Computing.* 34:227-231, 1996.

Bray, R. C., D. J. Butterwick, M. R. Doschak and J. V. Tyberg. Coloured microsphere assessment of blood flow to knee ligaments in adult rabbits: Effects of injury. *Journal Of Orthopaedic Research.* 14:618-625, 1996.

Bremer, H. J., A. Anninos and B. Schulz. Amino Acid Composition Of Food Products Used In the Treatment Of Patients With Disorders Of the Amino Acid and Protein Metabolism. *European Journal of Pediatrics.* 155:S 108-S 114, 1996.

Brown, W. E., M. G. Magno, P. D. Buckman, M. F. Di, D. R. Gale and J. D. Mannion. The coronary collateral circulation in normal goats. *J. Surg. Res.* 51:54-9, 1991.

Brown, W. R., D. M. Moody, D. A. Stump, D. D. Deal and R. L. Anderson. Dog model for cerebrovascular studies of the proximal-to-distal distribution of sequentially injected emboli. *Microvasc Res.* 50:105-12., 1995.

Brunston, R. L., W. K. Tao, A. Bidani, D. L. Traber and J. B. Zwischenberger. Organ blood flow during arteriovenous carbon dioxide removal. *Asaio Journal.* 43:M821-M824, 1997.

Brust, P., R. Bauer, G. Vorwieger, B. Walter, R. Bergmann, F. Fuchtner, J. Steinbach, U. Zwiener and B. Johannsen. Upregulation of the aromatic amino acid decarboxylase under neonatal asphyxia. *Neurobiology of Disease*. 6:131-139, 1999.

Buchwalder, L. F., M. Lin, T. J. McDonald and P. W. Nathanielsz. Fetal sheep adrenal blood flow responses to hypoxemia after splanchnicotomy using fluorescent microspheres. *Journal Of Applied Physiology*. 84:82-89, 1998.

Buckberg, G. D., J. C. Luck, D. B. Payne, J. I. E. Hoffman, J. P. Archie and D. E. Fixler. Some sources of error in measuring regional blood flow with radioactive microspheres. *J. Appl. Physiol.* 31:598-604, 1971.

Burchfield, D. J., A. Pena, A. J. M. Peters, R. M. Abrams and D. Phillips. Cocaine does not compromise cerebral or myocardial oxygen delivery in fetal sheep. *Reproduction Fertility And Development*. 8:383-389, 1996.

Caldarone, C. A., I. B. Krukenkamp, B. D. Misare and S. Levitsky. Perfusion deficits with retrograde warm blood cardioplegia. *Ann Thorac Surg*. 57:403-6., 1994.

Carlson, G. D., Y. Minato, A. Okada, C. D. Gorden, K. E. Warden, J. M. Barbeau, C. L. Biro, E. Bahniuk, H. H. Bohlman and J. C. Lamanna. Early time-dependent decompression for spinal cord injury: Vascular mechanisms of recovery. *JOURNAL OF NEUROTRAUMA*. 14:951-962, 1997.

Carlson, G. D., K. E. Warden, J. M. Barbeau, E. Bahniuk, K. L. KutinaNelson, C. L. Biro, H. H. Bohlman and J. C. LaManna. Viscoelastic relaxation and regional blood flow response to spinal cord compression and decompression. *Spine*. 22:1285-1291, 1997.

Cerda, S. E., C. Y. Tong, D. D. Deal and J. C. Eisenach. A physiologic assessment of intrathecal amitriptyline in sheep. *Anesthesiology*. 86:1094-1103, 1997.

Chase, P. B., K. B. Kern, A. B. Sanders, C. W. Otto and G. A. Ewy. Effects of graded doses of epinephrine on both noninvasive and invasive measures of myocardial perfusion and blood flow during cardiopulmonary resuscitation. *Crit Care Med*. 21:413-9., 1993.

Chien, G. L., C. G. Anselone, R. F. Davis and D. M. Van-Winkle. Fluorescent vs. radioactive microsphere measurement of regional myocardial blood flow. *Cardiovasc Res*. 30:405-12, 1995.

Chiou, G. C. and Y. J. Chen. Effects of D- and L-isomers of timolol on retinal and choroidal blood flow in ocular hypertensive rabbit eyes. *J Ocul Pharmacol*. 8:183-90., 1992.

Chiou, G. C. and Y. J. Chen. Improvement of ocular blood flow with dopamine antagonists on ocular-hypertensive rabbit eyes. *Chung Kuo Yao Li Hsueh Pao*. 13:481-4., 1992.

Cicutti, N. and K. Rakusan. The effect of hypoxia on capillary flow direction in the isolated perfused rat heart. *Can J Cardiol*. 10:367-73, 1994.

Cicutti, N., K. Rakusan and H. F. Downey. Colored microspheres reveal interarterial microvascular anastomoses in canine myocardium. *Basic Res Cardiol.* 87:400-9, 1992.

Cicutti, N., K. Rakusan and H. F. Downey. Coronary artery occlusion extends perfusion territory boundaries through microvascular collaterals. *Basic Res Cardiol.* 89:427-37., 1994.

Cioffi, G. A. and S. Orgul. Intraluminal corrosion casting and colored-microsphere techniques as means to assess optic nerve hemodynamics. *Ocular Blood Flow.* 226:1996.

Cochran, R. P., K. S. Kunzelman, C. R. Vocelka, H. Akimoto, R. Thomas, L. O. Soltow and B. D. Spiess. Perfluorocarbon emulsion in the cardiopulmonary bypass prime reduces neurologic injury. *Annals Of Thoracic Surgery.* 63:1326-1332, 1997.

Conhaim, R. L. and L. A. Rodenkirch. Estimated functional diameter of alveolar septal microvessels in zone 1. *Am J Physiol.* 271:H996-1003., 1996.

Consigny, P. M., E. D. Verrier, B. D. Payne, G. Edelist, J. Jester, R. W. Baer, G. J. Vlahakes and J. I. Hoffman. Acute and chronic microsphere loss from canine left ventricular myocardium. *Am J Physiol.* 242:H392-404, 1982.

Daldrup, H. E., D. M. Shames, W. Husseini, M. F. Wendland, Y. Okuhata and R. C. Brasch. Quantification of the extraction fraction for gadopentetate across breast cancer capillaries. *MAGNETIC RESONANCE IN MEDICINE.* 40:537-543, 1998.

DeBehnke, D. J., M. G. Angelos and J. E. Leasure. Use of cardiopulmonary bypass, high-dose epinephrine, and standard-dose epinephrine in resuscitation from post-countershock electromechanical dissociation. *Ann Emerg Med.* 21:1051-7., 1992.

Deem, S., M. J. Bishop and M. K. Alberts. Effect of anemia on intrapulmonary shunt during atelectasis in rabbits. *Journal Of Applied Physiology.* 79:1951-1957, 1995.

Degens, H., A. J. Craven, J. C. Jarvis and S. Salmons. The use of coloured dye-extraction microspheres to measure blood flow in rabbit skeletal muscle: A validation study with special emphasis on repeated measurements. *Experimental Physiology.* 81:239-249, 1996.

Deka, C., L. A. Sklar and J. A. Steinkamp. Fluorescence lifetime measurements in a flow cytometer by amplitude demodulation using digital data acquisition technique. *Cytometry.* 17:94-101, 1994.

Deveci, D. and S. Egginton. Development of the fluorescent microsphere technique for quantifying regional blood flow in small mammals. *EXPERIMENTAL-PHYSIOLOGY.* 84:615-630, 1999.

Diebel, L. N., J. G. Tyburski and S. A. Dulchavsky. Effect of hypertonic saline solution and dextran on ventricular blood flow and heart-lung interaction after hemorrhagic shock. *Surgery.* 124:642-650, 1998.

Domenech, R. J., J. I. Hoffman, M. I. Noble, K. B. Saunders, J. R. Henson and S. Subijanto. Total and regional coronary blood flow measured by radioactive microspheres in conscious and anesthetized dogs. *Circ Res.* 25:581-96, 1969.

- Eckstein, E. C., J. F. Koleski and C. M. Waters. Concentration profiles of 1 and 2.5 microns beads during blood flow. *Hematocrit effects. Asaio Trans.* 35:188-90, 1989.
- Ehring, T., D. Baumgart, M. Krajcar, M. Hummelgen, S. Kompa and G. Heusch. Attenuation of myocardial stunning by the ACE inhibitor ramiprilat through a signal cascade of bradykinin and prostaglandins but not nitric oxide. *Circulation.* 90:1368-85., 1994.
- Ehring, T., M. Bohm and G. Heusch. The calcium antagonist nisoldipine improves the functional recovery of reperfused myocardium only when given before ischemia. *J Cardiovasc Pharmacol.* 20:63-74., 1992.
- Eising, G. P., L. Mao, G. W. SchmidSchonbein, R. L. Engler and J. Ross. Effects of induced tolerance to bacterial lipopolysaccharide on myocardial infarct size in rats. *Cardiovascular Research.* 31:73-81, 1996.
- Elgjo, G. I., B. P. Mathew, L. F. P. deFigueiredo, P. J. Schenarts, J. W. Horton, M. A. Dubick and G. C. Kramer. Resuscitation with hypertonic saline dextran improves cardiac function in vivo and ex vivo after burn injury in sheep. *SHOCK.* 9:375-383, 1998.
- Erickson, H. H., S. L. Bernard, R. W. Glenny, M. R. Fedde, N. L. Polissar, R. J. Basaraba, S. M. Walther, E. M. Gaughan, R. McMurphy and M. P. Hlastala. Effect of furosemide on pulmonary blood flow distribution in resting and exercising horses. *Journal Of Applied Physiology.* 86:2034-2043, 1999.
- Fan, F. C., G. B. Schuessler, R. Y. Z. Chen and S. Chien. Determination of blood flow and shunting of 9- and 15- μm spheres in regional beds. *Am. J. Physiol.* 237:H25-H33, 1979.
- Fenton, B. M., R. T. Carr and G. R. Cokelet. Nonuniform red cell distribution in 20 to 100 micrometers bifurcations. *Microvasc Res.* 29:103-26, 1985.
- Flynn, A. E., D. L. Coggins, R. E. Austin, D. D. Muehrcke, G. S. Aldea, M. Goto, J. W. Doucette and J. I. Hoffman. Nonuniform blood flow in the canine left ventricle. *J Surg Res.* 49:379-84, 1990.
- Forrester, K., M. Doschak and R. Bray. In vivo comparison of scanning technique and wavelength in laser Doppler perfusion imaging: measurement in knee ligaments of adult rabbits. *MEDICAL & BIOLOGICAL ENGINEERING & COMPUTING.* 35:581-586, 1997.
- Frangioni, G. and G. Borgioli. Data on haemodynamics of splenic circulation in the newt. *Journal Of Zoology.* 238:149-155, 1996.
- Fujiki, H., T. Mori, K. Yoshida, T. Imaizumi and M. Tominaga. OPC-18790, a novel positive inotropic agent, has both arterial and venous vascular dilating actions in the dog. *European Journal Of Pharmacology.* 313:191-200, 1996.

Funato, H., M. Watanabe and A. Uemura. Therapeutic effects of a calcium antagonist, lacidipine, on stroke-prone spontaneously hypertensive rats with cerebrovascular lesions. *JAPANESE-JOURNAL-OF-PHARMACOLOGY*. 80:199-208, 1999.

Gagnon, R., J. Challis, L. Johnston and L. Fraher. Fetal endocrine responses to chronic placental embolization in the late-gestation ovine fetus. *Am J Obstet Gynecol.* 170:929-38., 1994.

Galiuto, L., A. N. DeMaria, K. MayNewman, U. DelBalzo, K. Ohmori, V. Bhargava, S. F. Flaim and S. Iliceto. Evaluation of dynamic changes in microvascular flow during ischemia-reperfusion by myocardial contrast echocardiography. *Journal of the American College of Cardiology*. 32:1096-1101, 1998.

Gates, R. N., J. G. Lee, H. Laks, D. C. Drinkwater, E. Rhudis, A. S. Aharon, J. Y. Chung and P. A. Chang. Evidence of improved microvascular perfusion when using antegrade and retrograde cardioplegia. *Annals Of Thoracic Surgery*. 62:1388-1391, 1996.

Gervais, M., P. Demolis, V. Domergue, M. Lesage, C. Richer and J. F. Giudicelli. Systemic and regional hemodynamics assessment in rats with fluorescent microspheres. *Journal of Cardiovascular Pharmacology*. 33:425-432, 1999.

Glenny, R. W. Blood flow measurements using fluorescent microspheres. Second Oxford Conference on Spectrometry. 1995.

Glenny, R. W., S. Bernard and M. Brinkley. Validation of fluorescent-labeled microspheres for measurement of regional organ perfusion. *J Appl Physiol.* 74:2585-97, 1993.

Glenny, R. W., S. Bernard, H. T. Robertson and M. P. Hlastala. Gravity is an important but secondary determinant of regional pulmonary blood flow in upright primates. *Journal of Applied Physiology*. 86:623-632, 1999.

Glenny, R. W., S. McKinney and K. T. Robertson. Spatial pattern of pulmonary blood flow distribution is stable over days. *Journal Of Applied Physiology*. 82:902-907, 1997.

Glenny, R. W., N. L. Polissar, S. McKinney and H. T. Robertson. Temporal heterogeneity of regional pulmonary perfusion is spatially clustered. *J Appl Physiol.* 79:986-1001., 1995.

Glenny, R. W. and H. T. Robertson. A computer simulation of pulmonary perfusion in three dimensions. *J Appl Physiol.* 79:357-69., 1995.

Gottlieb, G. J., S. H. Kubo and D. R. Alonso. Ultrastructural characterization of the border zone surrounding early experimental myocardial infarcts in dogs. *Am J Pathol.* 103:292-303, 1981.

Granetzny, A., U. Schwanke, C. Schmitz, G. Arnold, D. Schafer, H. D. Schulte, E. Gams and J. D. Schipke. Pharmacologic heart rate reduction: Effect of a novel, specific bradycardic agent on the heart. *THORACIC AND CARDIOVASCULAR SURGEON*. 46:63-69, 1998.

- Granetzny, A., U. Schwanke, C. Schmitz, S. SchmitzSpanke, G. Arnold, H. D. Schulte and J. D. Schipke. Effects of a new bradycardic agent on the isolated rabbit heart. *Zeitschrift Fur Kardiologie*. 85:953-960, 1996.
- Greve, G. and T. Saetersdal. Problems related to infarct size measurements in the rat heart. *Acta Anat Basel*. 142:366-73, 1991.
- Groban, L., D. A. Zvara, D. D. Deal, J. C. Vernon, C. W. Flye, X. L. Ma and J. VintenJohansen. Cloricromene reduces infarct size and alters postischaemic blood flow defects in dog myocardium. *CLINICAL AND EXPERIMENTAL PHARMACOLOGY AND PHYSIOLOGY*. 25:417-423, 1998.
- Gross, T. S., A. A. Damji, S. Judex, R. C. Bray and R. F. Zernicke. Bone hyperemia precedes disuse-induced intracortical bone resorption. *Journal of Applied Physiology*. 86:230-235, 1999.
- Gross, W., P. Zeller, U. Kreimeier and K. Messmer. Pseudocolor display of regional organ blood flow determined by means of the radioactive microsphere technique. *Comput Med Imaging Graph*. 14:1-11, 1990.
- Grund, F., H. T. Sommerschild, T. Lyberg, K. A. Kirkeboen and A. Ilebekk. Microembolization in pigs: effects on coronary blood flow and myocardial ischemic tolerance. *AMERICAN-JOURNAL-OF-PHYSIOLOGY-HEART-AND-CIRCULATORY-PHYSIOLOGY*. 46:H533-H542, 1999.
- Guilbault, G. G. Practical Fluorescence. Modern Monographs in Analytical Chemistry. 3: 1990.
- Haga, Y., R. Nordlander, P. O. Sjoquist and L. Ryd'en. Influence of coronary venous retroinfusion and vasodilatation on regional myocardial blood flow measurement with microspheres. An analysis of 'microsphere loss' from ischaemic and reperfused porcine hearts. *Acta Physiol Scand*. 153:13-20, 1995.
- Hagendorff, A., C. Dettmers, P. Danos, M. Hummelgen, C. Vahlhaus, C. Martin, G. Heusch and B. Luderitz. Cerebral vasoconstriction during sustained ventricular tachycardia induces an ischemic stress response of brain tissue in rats. *Journal of Molecular and Cellular Cardiology*. 30:2081-2094, 1998.
- Hagendorff, A., C. Vahlhaus, W. Jung, C. Martin, G. Heusch and B. Luderitz. Association between hemodynamic parameters and the degeneration of sustained ventricular tachycardias into ventricular fibrillation in rats. *JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY*. 29:3091-3103, 1997.
- Hakkinen, J. P., M. W. Miller, A. H. Smith and D. R. Knight. Measurement of organ blood flow with coloured microspheres in the rat. *Cardiovasc Res*. 29:74-9, 1995.
- Hale, S. L., K. J. Alker and R. A. Kloner. Evaluation of nonradioactive, colored microspheres for measurement of regional myocardial blood flow in dogs. *Circulation*. 78:428-34, 1988.
- Hale, S. L. and R. A. Kloner. Effect of early coronary artery reperfusion on infarct development in a model of low collateral flow. *Cardiovasc Res*. 21:668-73, 1987.

Hale, S. L., M. T. Vivaldi and R. A. Kloner. Fluorescent microspheres: a new tool for visualization of ischemic myocardium in rats. *Am J Physiol.* 251:H863-8, 1986.

Hales, J. R. S. and W. J. Cliff. Direct observation of the behavior of microspheres in microvasculature. *Bibl. Anat.* 15:87-91, 1977.

Hall, J. L., L. A. Hernandez, J. Henderson, L. A. Kellerman and W. C. Stanley. Decreased interstitial glucose and transmural gradient in lactate during ischemia. *Basic Res Cardiol.* 89:468-86, 1994.

Hanger, C. C., S. C. Hillier, R. G. Presson, Jr., R. W. Glenny and W. W. Wagner, Jr. Measuring pulmonary microvessel diameters using video image analysis. *J Appl Physiol.* 79:526-32., 1995.

Hardy, J., A. L. Bertone and W. W. Muir. Joint pressure influences synovial tissue blood flow as determined by colored microspheres. *Journal Of Applied Physiology.* 80:1225-1232, 1996.

Hariawala, M. D., J. R. Horowitz, D. Esakof, D. D. Sheriff, D. Walter, B. Keyt, J. M. Isner and J. F. Symes. VEGF improves myocardial blood flow but produces EDRF-mediated hypotension in porcine hearts. *Journal Of Surgical Research.* 63:77-82, 1996.

Harris, D. A. Spectrophotometry and Spectrofluorimetry a Practical Approach, Washington, D.C.: IRL Press, 1987

Hasegawa, T., A. Kimura, R. Miyataka, M. Inagaki and K. Ishikawa. Basic fibroblast growth factor increases regional myocardial blood flow and salvages myocardium in the infarct border zone in a rabbit model of acute myocardial infarction. *Angiology.* 50:487-495, 1999.

Hatori, N., Y. Uriuda, K. Isozima, T. Isono, E. Okuda, K. Hamada, I. Nakahoshi, A. Kurita, H. Yoshizu and S. Tanaka. Short-term treatment with synchronized coronary venous retroperfusion before full reperfusion significantly reduces myocardial infarct size. *Am Heart J.* 123:1166-74., 1992.

Herijgers, P., V. Leunens, T. B. Tjandramaga, K. Mubagwa and W. Flameng. Changes in organ perfusion after brain death in the rat and its relation to circulating catecholamines. *Transplantation.* 62:330-335, 1996.

Heyman, M. A., B. D. Payne, J. I. Hoffman and A. M. Rudolf. Blood flow measurements with radionuclide-labeled particles. *Prog. Cardiovasc. Dis.* 20:55-79, 1977.

Hiller, K. H., P. Adami, S. Voll, F. Roder, P. Kowallik, W. R. Bauer, A. Haase and G. Ertl. In vivo colored microspheres in the isolated rat heart for use in NMR. *Journal Of Molecular And Cellular Cardiology.* 28:571-577, 1996.

Hiller, K. H., F. Roder, P. Adami, S. Voll, P. Kowallik, A. Haase, G. Ertl and W. R. Bauer. Study of microcirculation by coloured microspheres and NMR-microscopy in isolated rat heart: Effect of ischaemia, endothelin-1 and endothelin-1 antagonist BQ 610. *JOURNAL OF MOLECULAR AND CELLULAR CARDIOLOGY.* 29:3115-3122, 1997.

Hillier, S. C., J. A. Graham, C. C. Hanger, P. S. Godbey, R. W. Glenny and W. W. Wagner. Hypoxic vasoconstriction in pulmonary arterioles and venules. *Journal Of Applied Physiology*. 82:1084-1090, 1997.

Hlastala, M. P., S. L. Bernard, H. H. Erickson, M. R. Fedde, E. M. Gaughan, R. McMurphy, M. J. Emery, N. Polissar and R. W. Glenny. Pulmonary blood flow distribution in standing horses is not dominated by gravity. *Journal Of Applied Physiology*. 81:1051-1061, 1996.

Hlastala, M. P., M. A. Chornuk, D. A. Self, H. J. Kallas, J. W. Burns, S. Bernard, N. L. Polissar and R. W. Glenny. Pulmonary blood flow redistribution by increased gravitational force. *JOURNAL OF APPLIED PHYSIOLOGY*. 84:1278-1288, 1998.

Hodeige, D., P.-M. De, W. Eechaute, J. Weyne and G. R. Heyndrickx. On the validity of blood flow measurement using colored microspheres. *American Journal of Physiology-Heart and Circulatory Physiology*. 276:H1150-H1158, 1999.

Hoffbrand, B. I. and R. P. Forsyth. Validity studies of the radioactive microsphere method for the study of the distribution of cardiac output, organ blood flow, and resistance in the conscious rhesus monkey. *Cardiovasc. Res.* 3:426-432, 1969.

Hong, S. J., K. Y. Wu and I. J. Chen. Ocular hypotensive and vasodilative effects of two beta-adrenergic blockers with intrinsic sympathomimetic activity. *CURRENT EYE RESEARCH*. 17:700-707, 1998.

Hood, D. D., J. C. Eisenach, C. Tong, E. Tommasi and T. L. Yaksh. Cardiorespiratory and spinal cord blood flow effects of intrathecal neostigmine methylsulfate, clonidine, and their combination in sheep [see comments]. *Anesthesiology*. 82:428-35., 1995.

Hooper, S. B. Role of luminal volume changes in the increase in pulmonary blood flow at birth in sheep. *Experimental Physiology*. 83:833-842, 1998.

Hoshida, S., T. Kuzuya, M. Nishida, N. Yamashita, H. Oe, M. Hori, T. Kamada and M. Tada. Adenosine blockade during reperfusion reverses the infarct limiting effect in preconditioned canine hearts. *Cardiovasc Res.* 28:1083-8, 1994.

Hoshida, S., T. Kuzuya, N. Yamashita, M. Nishida, S. Kitahara, M. Hori, T. Kamada and M. Tada. gamma-Glutamylcysteine ethyl ester for myocardial protection in dogs during ischemia and reperfusion. *J Am Coll Cardiol.* 1. 24:1391-7., 1994.

Hu, N., T. D. Ngo and E. B. Clark. Distribution of blood flow between embryo and vitelline bed in the stage 18, 21 and 24 chick embryo. *Cardiovascular Research*. 31:131, 1996.

Huijbregts, P., E. J. M. Feskens, L. Rasanen, A. Albertifidanza, M. Mutanen, F. Fidanza and D. Kromhout. Dietary Intake In Five Ageing Cohorts Of Men In Finland, Italy and the Netherlands. *European Journal of Clinical Nutrition*. 49:852-860, 1995.

Iglesiasbarreira, V., M. T. Ahn, B. Reusens, S. Dahri, J. J. Hoet and C. Remacle. Pre- and postnatal low protein diet affect pancreatic islet blood flow and insulin release in adult rats. *Endocrinology*. 137:3797-3801, 1996.

Ikeda, T., Y. Murata, E. J. Quilligan, P. Cifuentes, S. Doi and S. D. Park. Two sinusoidal heart rate patterns in fetal lambs undergoing extracorporeal membrane oxygenation. *American Journal of Obstetrics and Gynecology*. 180:462-468, 1999.

Inada, Y., S. Tamai, S. Mizumoto, H. Ono, K. Kawanishi and A. Fukui. Non-radioactive coloured microsphere measurement of regional tissue blood flow for axial pattern flaps in rabbits. *Br J Plast Surg*. 46:127-31, 1993.

Ishikawa, K., N. Kamata, S. Nakai, H. Akiyama, H. Koka, I. Ogawa and R. Katori. Preservation of high regional blood flow at epicardial rim after coronary occlusion in dogs. *Am J Physiol*. 267:H528-34, 1994.

Ishikawa, K., I. Ogawa, M. Shimizu, H. Koka, N. Kamata, S. Nakai, H. Akiyama, T. Shimamoto, N. Ishida and K. Kanamasa. The importance of good endocardial reflow immediately after reperfusion for myocardial salvage in dogs. *Jpn. Circ. J.* 55:983-93, 1991.

Ishikawa, K., I. Ogawa, M. Shimizu, H. Koka, N. Kamata, S. Nakai and R. Katori. Residual critical coronary stenosis during myocardial reperfusion is deleterious to myocardial salvage in dogs. *Jpn Circ J*. 56:921-8., 1992.

Ishikawa, M., T. Mori, S. Itoh, H. Fujiki, K. Koga, M. Tominaga and Y. Yabuuchi. Effects of OPC-18790, a new positive inotropic agent, on energetics in the ischaemic canine heart: a 31P-MRS study. *Cardiovasc Res*. 30:299-306., 1995.

Jasper, M. S., P. McDermott, D. S. Gann and W. C. Engeland. Measurement of blood flow to the adrenal capsule, cortex and medulla in dogs after hemorrhage by fluorescent microspheres. *J Auton Nerv Syst*. 30:159-67, 1990.

Johansen, B., M. N. Melsom, T. Flatebo and G. Nicolaysen. Time course and pattern of pulmonary flow distribution following unilateral airway occlusion in sheep. *CLINICAL SCIENCE*. 94:453-460, 1998.

Kaihara, S., P. D. Heerden, T. Migita and H. N. Wagner Jr. Measurement of distribution of cardiac output. *Journal of Applied Physiology*. 25:696-700, 1968.

Kallas, H. J., K. B. Domino, R. W. Glenny, E. A. Anderson and M. P. Hlastala. Pulmonary blood flow redistribution with low levels of positive end-expiratory pressure. *ANESTHESIOLOGY*. 88:1291-1299, 1998.

Kamimura, T., H. Sakamoto and K. Misumi. Regional blood flow distribution from the proximal arterial cannula during veno-arterial extracorporeal membrane oxygenation in neonatal dog. *Journal of Veterinary Medical Science*. 61:311-315, 1999.

Kanagawa, R., T. Wada, T. Sanada, M. Ojima and Y. Inada. Regional hemodynamic effects of candesartan cilexetil (TCV-116), an angiotensin II AT(1)-receptor antagonist, in conscious spontaneously hypertensive rats. *Japanese Journal Of Pharmacology.* 73:185-190, 1997.

Kapadia, S. J., J. S. Terlato and A. S. Most. Presence of a critical coronary artery stenosis does not abolish the protective effect of ischemic preconditioning. *Circulation.* 95:1286-1292, 1997.

Kawagoe, Y., S. Permutt and H. E. Fessler. Hyperinflation with intrinsic PEEP and respiratory muscle blood flow. *J Appl Physiol.* 77:2440-8., 1994.

Kelly, R. F., T. L. Hursey, R. B. Patel, J. E. Parrillo and G. L. Schaer. Effect of poloxamer 188 on collateral blood flow, myocardial infarct size, and left ventricular function in a canine model of prolonged (3-hour) coronary occlusion and reperfusion. *JOURNAL OF THROMBOSIS AND THROMBOLYSIS.* 5:239-247, 1998.

Kelly, R. F., T. L. Hursey, G. L. Schaer, M. J. Piotrowski, S. V. Dee, J. E. Parrillo and S. M. Hollenberg. Cardiac endothelin release and infarct size, myocardial blood flow, and ventricular function in canine infarction and reperfusion. *Journal Of Investigative Medicine.* 44:575-582, 1996.

Kemp, P. A., S. M. Gardiner, J. E. March, P. C. Rubin and T. Bennett. Assessment of the effects of endothelin-1 and magnesium sulphate on regional blood flows in conscious rats, by the coloured microsphere reference technique. *British Journal of Pharmacology.* 126:621-626, 1999.

Kern, K. B. and G. A. Ewy. Minimal coronary stenoses and left ventricular blood flow during CPR [published erratum appears in Ann Emerg Med 1993 Jan;22(1):124]. *Ann Emerg Med.* 21:1066-72., 1992.

Kern, K. B., L. Lancaster, S. Goldman and G. A. Ewy. The effect of coronary artery lesions on the relationship between coronary perfusion pressure and myocardial blood flow during cardiopulmonary resuscitation in pigs. *Am. Heart J.* 120:324-33, 1990.

Khoobehi, B., B. Shoelson, Y. Z. Zhang and G. A. Peyman. Fluorescent microsphere imaging: A particle-tracking approach to the hemodynamic assessment of the retina and choroid. *Ophthalmic Surgery And Lasers.* 28:937-947, 1997.

Kimura, A., K. Ishikawa and I. Ogawa. Myocardial salvage by reperfusion 12 hours after coronary ligation in dogs. *JAPANESE CIRCULATION JOURNAL-ENGLISH EDITION.* 62:294-298, 1998.

Kinoshita, H., I. Ijiri, S. Ameno, C. Fuke, Y. Fujisawa and K. Ameno. Inhibitory mechanism of intestinal ethanol absorption induced by high acetaldehyde concentrations: Effect of intestinal blood flow and substance specificity. *Alcoholism Clinical And Experimental Research.* 20:510-513, 1996.

Kinoshita, K., D. J. Hearse, M. V. Braimbridge and A. S. Manning. 'Early' ischemia and reperfusion-induced arrhythmias: antiarrhythmic effects of diltiazem in the conscious rat. *Can J Cardiol.* 4:37-43, 1988.

Klabunde, R. E., W. A. Anderson, M. Locke, S. E. Ianuzzo and C. D. Ianuzzo. Regional blood flows in the goat latissimus dorsi muscle before and after chronic stimulation. *Journal Of Applied Physiology*. 81:2365-2372, 1996.

Klemm, K. and F. G. Moody. Regional intestinal blood flow and nitric oxide synthase inhibition during sepsis in the rat. *Annals of Surgery*. 227:126-133, 1998.

Kobayashi, N., K. Kobayashi, K. Hara, T. Higashi, H. Yanaka, S. Yagi and H. Matsuoka. Benidipine stimulates nitric oxide synthase and improves coronary circulation in hypertensive rats. *American Journal of Hypertension*. 12:483-491, 1999.

Kobayashi, N., K. Kobayashi, K. Kono, T. Akabane, H. Kaneko, M. Takada, N. Tsuchiya and S. Yagi. Effects of microsphere suspension agents on systemic hemodynamics in rats. Comparison of nonradioactive colored and radioactive microspheres. *Jpn Heart J*. 35:467-75, 1994.

Kobayashi, N., K. Kobayashi, K. Kouno, S. Horinaka and S. Yagi. Effects of intra-atrial injection of colored microspheres on systemic hemodynamics and regional blood flow in rats. *Am J Physiol*. 266:H1910-7., 1994.

Kobayashi, N., K. Kobayashi, K. Kouno, S. Yagi and H. Matsuoka. Effect of benidipine on microvascular remodeling and coronary flow reserve in two-kidney, one clip Goldblatt hypertension. *Journal Of Hypertension*. 15:1285-1294, 1997.

Kohmoto, T., P. E. Fisher, A. Gu, S. M. Zhu, C. M. DeRosa, C. R. Smith and D. Burkhoff. Physiology, histology, and 2-week morphology of acute transmyocardial channels made with a CO₂ laser. *Annals Of Thoracic Surgery*. 63:1275-1283, 1997.

Kohmoto, T., P. E. Fisher, A. Gu, S. M. Zhu, O. J. Yano, H. M. Spotnitz, C. R. Smith and D. Burkhoff. Does blood flow through holmium:YAG transmyocardial laser channels? *Annals Of Thoracic Surgery*. 61:861-868, 1996.

Kohmoto, T., G. Uzun, A. Gu, S. M. Zhu, C. R. Smith and D. Burkhoff. Blood flow capacity via direct acute myocardial revascularization. *Basic Research In Cardiology*. 92:45-51, 1997.

Kowallik, P., D. Baumgart, A. Skyschally, T. Ehring and G. Heusch. Three-dimensional analysis of regional mechanical function, blood flow and electrophysiological parameters during early myocardial ischemia in dogs. *Basic Res Cardiol*. 87:215-26., 1992.

Kowallik, P., R. Schulz, B. D. Guth, A. Schade, W. Paffhausen, R. Gross and G. Heusch. Measurement of regional myocardial blood flow with multiple colored microspheres. *Circulation*. 83:974-82, 1991.

Kramer, C. M., P. D. Nicol, W. J. Rogers, M. M. Suzuki, A. Shaffer, T. M. Theobald and N. Reichek. Reduced sympathetic innervation underlies adjacent noninfarcted region dysfunction during left ventricular remodeling. *Journal Of The American College Of Cardiology*. 30:1079-1085, 1997.

Kreidstein, M. L., R. H. Levine, R. J. Knowlton and C. Y. Pang. Serial fluorometric assessments of skin perfusion in isolated perfused human skin flaps. *Br J Plast Surg*. 48:288-93, 1995.

Kuhle, W. G., G. Porenta, S. C. Huang, D. Buxton, S. S. Gambhir, H. Hansen, M. E. Phelps and H. R. Schelbert. Quantification of regional myocardial blood flow using ¹³N-ammonia and reoriented dynamic positron emission tomographic imaging. *Circulation*. 86:1004-17, 1992.

Kurdak, S. S., B. Grassi, P. D. Wagner and M. C. Hogan. Blood flow distribution in working in situ canine muscle during blood flow reduction. *Journal Of Applied Physiology*. 80:1978-1983, 1996.

Kuwahira, I., N. C. Gonzalez, N. Heisler and J. Piiper. Regional blood flow in conscious resting rats determined by microsphere distribution. *J Appl Physiol*. 74:203-10, 1993.

Kuwahira, I., H. Mori, Y. Moue, Y. Shinozaki, Y. Ohta, H. Yamabayashi, H. Okino, N. C. Gonzalez, N. Heisler and J. Piiper. Cardiac output and regional blood flow measurement with nonradioactive microspheres by X-ray fluorescence spectrometry in rats. *Adv Exp Med Biol*. 345:877-84, 1994.

Kuwahira, I., Y. Moue, Y. Ohta and N. C. Gonzalez. Chronic hypoxia decreases heterogeneity of pulmonary blood flow distribution in rats. *Respiration Physiology*. 104:2-3, 1996.

Kvinnslund, S., K. Heyeraas and E. S. Ofjord. Effect of experimental tooth movement on periodontal and pulpal blood flow. *Eur J Orthod*. 11:200-5, 1989.

Kvinnslund, S., A. B. Kristiansen, I. Kvinnslund and K. J. Heyeraas. Effect of experimental traumatic occlusion on periodontal and pulpal blood flow. *Acta Odontol Scand*. 50:211-9, 1992.

Kvinnslund, S., I. Kvinnslund and A. B. Kristiansen. Effect of experimental traumatic occlusion on blood flow in the temporomandibular joint of the rat. *Acta Odontol Scand*. 51:293-8, 1993.

Laham, R. J., M. Simons, M. Tofukuji, D. Hung and F. W. Sellke. Modulation of myocardial perfusion and vascular reactivity by pericardial basic fibroblast growth factor: Insight into ischemia-induced reduction in endothelium-dependent vasodilatation. *Journal of Thoracic and Cardiovascular Surgery*. 116:1022-1028, 1998.

Lee, J., R. N. Gates, H. Laks, D. C. Drinkwater, E. Rhudis, A. Aharon, A. Ardehali and P. Chang. A comparison of distribution between simultaneously or sequentially delivered antegrade/retrograde blood cardioplegia. *Journal Of Cardiac Surgery*. 11:111-115, 1996.

Li, D. S., A. C. Yong and D. Kilpatrick. Validation of a subendocardial ischaemic sheep model by intracoronary fluorescent microspheres. *Clinical & Experimental Pharmacology & Physiology*. 23:111-118, 1996.

Liu, S. X. L., C. H. Chiang, Q. S. Yao and G. C. Y. Chiou. Increase of ocular blood flow by some phytogenic compounds. *Journal Of Ocular Pharmacology And Therapeutics*. 12:95-101, 1996.

Liu, Y. H., R. C. Bahn and E. L. Ritman. Microvascular blood volume-to-flow relationships in porcine heart wall: whole body CT evaluation in vivo. *Am J Physiol*. 269:H1820-6., 1995.

Liu, Y. M., P. H. Guth, K. Kaneko, E. H. Livingston and F. C. Brunicardi. Dynamic in vivo observation of rat islet microcirculation. *Pancreas*. 8:15-21, 1993.

Lossec, G., C. Duchamp, Y. Lebreton and P. Herpin-. Postnatal changes in regional blood flow during cold-induced shivering in sow-reared piglets. *CANADIAN-JOURNAL-OF-PHYSIOLOGY-AND-PHARMACOLOGY*. 77:414-421, 1999.

Luchtel, D. L., J. C. Boykin, S. L. Bernard and R. W. Glenny. Histological methods to determine blood flow distribution with fluorescent microspheres. *Biotech Histochem*. 73:291-309, 1998.

Mahgoub, M. A., J. H. Guo, S. P. Gao, M. M. Taher, D. D. Salter, A. S. Wechsler and A. S. Abd-Elfattah. Hyperdynamic circulation of arteriovenous fistula preconditions the heart and limits infarct size. *ANNALS-OF-THORACIC-SURGERY*. 68:22-28, 1999.

Main, M. L., J. F. Escobar, S. A. Hall, A. L. Killam and P. A. Grayburn. Detection of myocardial perfusion defects by contrast echocardiography in the setting of acute myocardial ischemia with residual antegrade flow. *JOURNAL OF THE AMERICAN SOCIETY OF ECHOCARDIOGRAPHY*. 11:228-235, 1998.

Mann, C. M., K. B. Domino, S. M. Walther, R. W. Glenny, N. L. Polissar and M. P. Hlastala. Redistribution of pulmonary blood flow during unilateral hypoxia in prone and supine dogs. *JOURNAL OF APPLIED PHYSIOLOGY*. 84:2010-2019, 1998.

Mannion, J. D., P. D. Buckman, M. G. Magno and F. Dimeo. Collateral blood flow from skeletal muscle to normal myocardium. *J Surg Res*. 53:578-87., 1992.

Mannion, J. D., M. G. Magno, P. D. Buckman, F. DiMeo, R. Greene, M. Bowers, M. McHugh and H. Menduke. Acute electrical stimulation increases extramyocardial collateral blood flow after a cardiomyoplasty. *Ann Thorac Surg*. 56:1351-8., 1993.

Marber, M. S., D. M. Walker, D. J. Eveson, J. M. Walker and D. M. Yellon. A single five minute period of rapid atrial pacing fails to limit infarct size in the in situ rabbit heart. *Cardiovasc Res*. 27:597-601, 1993.

Marshall, W. G., G. B. Boatman, G. Dickerson, A. Perlin, E. P. Todd and J. R. Utley. Shunting, release, and distribution of nine and fifteen micron spheres in myocardium. *Surgery*. 79:631-7, 1976.

Masuda, Y., M. Ozaki and T. Oguma. Alteration of hepatic microcirculation by oxethazaine and some vasoconstrictors in the perfused rat liver. *Biochemical Pharmacology*. 53:1779-1787, 1997.

Maxwell, A. J., E. Schable, D. Bernstein and J. P. Cooke. Limb blood flow during exercise is dependent on nitric oxide. *CIRCULATION*. 98:369-374, 1998.

Maxwell, L. C., A. P. Shepherd, G. L. Riedel and M. D. Morris. Effect of microsphere size on apparent intramural distribution of intestinal blood flow. *Am J Physiol*. 241:H408-14, 1981.

Mazhari, R., J. H. Omens, L. K. Waldman and A. D. McCulloch. Regional myocardial perfusion and mechanics: A model-based method of analysis. *ANNALS OF BIOMEDICAL ENGINEERING*. 26:743-755, 1998.

Mazoit, J. X., R. LeGuen, A. Decaux, P. Albaladejo and K. Samii. Application of HPLC to counting of colored microspheres in determination of regional blood flow-Special Communication. *AMERICAN JOURNAL OF PHYSIOLOGY-HEART AND CIRCULATORY PHYSIOLOGY*. 45:H1041-H1047, 1998.

McCraib, G. J. and R. Harding. Cerebral blood flow is increased throughout 12 h of hypoxaemia in the mid-gestation ovine fetus. *Reprod Fertil Dev*. 7:463-7, 1995.

McCraib, G. J. and R. Harding. Role of nitric oxide in the regulation of cerebral blood flow in the ovine foetus. *Clinical And Experimental Pharmacology And Physiology*. 23:10-11, 1996.

McCraib, G. J. and R. Harding. Cerebral oxygen delivery is reduced during the acidaemia associated with prolonged hypoxaemia in the immature ovine fetus. *Biology Of The Neonate*. 71:385-394, 1997.

McCulloch, A. D., D. Sung, J. M. Wilson, R. S. Pavelec and J. H. Omens. Flow-function relations during graded coronary occlusions in the dog: effects of transmural location and segment orientation. *CARDIOVASCULAR RESEARCH*. 37:636-645, 1998.

McHugh, N. A., A. Solowiej, R. E. Klabunde and G. F. Merrill. Acute coronary vascular and myocardial perfusion effects of conjugated equine estrogen in the anesthetized dog. *Basic Research in Cardiology*. 93:470-476, 1998.

McMahan, C. A., L. C. Maxwell and A. P. Shepherd. Estimation of the distribution of blood vessel diameters from the arteriovenous passage of microspheres. *Biometrics*. 42:371-80, 1986.

Medvedev, O. S., E. R. Martynova, R. S. Akchurin and V. Y. Khalatov. Experimental estimation of chronic microsphere loss from the rat myocardium. *Biull. Eksp. Biol. Med.* 103:8-10, 1987.

Meissner, A., T. P. Weber, H. VanAken, K. Zbieranek and N. Rolf. Clonidine improves recovery from myocardial stunning in conscious chronically instrumented dogs. *Anesthesia and Analgesia*. 87:1009-1014, 1998.

Melsom, M. N., T. Flatebo and G. Nicolaysen. Hypoxia and hyperoxia both transiently affect distribution of pulmonary perfusion but not ventilation in awake sheep. *ACTA-PHYSIOLOGICA-SCANDINAVICA*. 166:151-158, 1999.

Melsom, M. N., T. Flatebo, O. V. Sjaastad, A. Aulie and G. Nicolaysen. Minor redistribution of ventilation and perfusion within the lung during exercise in sheep [In Process Citation]. *Acta Physiol Scand.* 165:283-92, 1999.

Melsom, M. N., J. Kramer-Johansen, T. Flatebo, C. Muller and G. Nicolaysen. Distribution of pulmonary ventilation and perfusion measured simultaneously in awake goats [published erratum appears in Acta Physiol Scand 1997 Jul;160(3):297]. *Acta Physiol Scand.* 159:199-208, 1997.

Mercer, D. W., K. Klemm, J. M. Cross, G. S. Smith, M. Cashman and T. A. Miller. Cholecystokinin-induced protection against gastric injury is independent of endogenous somatostatin. *American Journal Of Physiology Gastrointestinal And Liver Physiology*. 34:G692-G700, 1996.

Meyns, B., Y. Nishimura, R. Racz, R. Jashari and W. Flameng. Organ perfusion with hemopump device assistance with and without intraaortic balloon pumping. *Journal Of Thoracic And Cardiovascular Surgery*. 114:243-253, 1997.

Miglierina, R. Some applications of microspheres in flow cytometry. *Biol Cell*. 58:127-30, 1986.

Miller, S. L., K. Dickson, G. Jenkin and D. W. Walker. Physiological evidence for arteriovenous anastomoses in the uterine circulation of late-pregnant ewes. *CLINICAL AND EXPERIMENTAL PHARMACOLOGY AND PHYSIOLOGY*. 25:92-98, 1998.

Mitsumaru, A., R. Yozu, Y. Inoue, S. Tanaka, H. Yoshito, K. Kanda, Y. Tsutsui, N. Tsutsui and S. Kawada. Experimental study of combination of extraaortic balloon counterpulsation and ventricular assist cup to acute heart failure in dogs. *Asaio Journal*. 43:187-192, 1997.

Miyamae, M., H. Fujiwara, M. Kida, R. Yokota, M. Tanaka, M. Katsuragawa, K. Hasegawa, M. Ohura, K. Koga, Y. Yabuuchi and et al. Preconditioning improves energy metabolism during reperfusion but does not attenuate myocardial stunning in porcine hearts. *Circulation*. 88:223-34., 1993.

Miyataka, M., K. Ishikawa and R. Katori. Basic fibroblast growth factor increased regional myocardial blood flow and limited infarct size of acutely infarcted myocardium in dogs. *ANGIOLOGY*. 49:381-390, 1998.

Moore, R. M., J. Hardy and W. W. Muir. Mural blood flow distribution in the large colon of horses during low-flow ischemia and reperfusion. *Am J Vet Res*. 56:812-8, 1995.

Mor-Avi, V., D. David, S. Akselrod, Y. Bitton and I. Choshniak. Myocardial regional blood flow: quantitative measurement by computer analysis of contrast enhanced echocardiographic images. *Ultrasound Med Biol*. 19:619-33, 1993.

Mori, H., M. Chujo, S. Haruyama, H. Sakamoto, Y. Shinozaki, M. Uddin-Mohammed, A. Iida and H. Nakazawa. Local continuity of myocardial blood flow studied by monochromatic synchrotron radiation-excited x-ray fluorescence spectrometry. *Circ Res*. 76:1088-100, 1995.

Mori, H., S. Haruyama, Y. Shinozaki, H. Okino, A. Iida, R. Takanashi, I. Sakuma, W. K. Husseini, B. D. Payne and J. I. Hoffman. New nonradioactive microspheres and more sensitive X-ray fluorescence to measure regional blood flow. *Am J Physiol*. 263:H1946-57, 1992.

Mori, H., S. Haruyama, Y. Shinozaki, H. Sakamoto, M. Chujyo and I. Kuwahira. [Practice of the flow measurement using nonradioactive microspheres and X-ray fluorescence spectrometer]. *Kokyu To Junkan*. 41:447-50, 1993.

- Mori, H., S. Haruyama, Y. Shinozaki, H. Sakamoto, H. Okino, R. Takanashi and I. Sakuma. [Regional blood flow measurement with nonradioactive microspheres]. *Kokyu To Junkan*. 40:957-61., 1992.
- Morita, Y., B. D. Payne, G. S. Aldea, C. McWatters, W. Husseini, H. Mori, J. I. Hoffman and L. Kaufman. Local blood flow measured by fluorescence excitation of nonradioactive microspheres. *Am J Physiol*. 258:H1573-84, 1990.
- Moros, E. G., A. W. Dutton, R. B. Roemer, M. Burton and K. Hyynnen. Experimental evaluation of two simple thermal models using hyperthermia in muscle in vivo. *Int J Hyperthermia*. 9:581-98., 1993.
- Mortola, J. P., D. Merazzi and L. Naso. Blood flow to the brown adipose tissue of conscious young rabbits during hypoxia in cold and warm conditions. *Pflugers Archiv-European Journal of Physiology*. 437:255-260, 1999.
- Mulder, A. L. M., J. C. vanGolde, F. W. Prinzen and C. E. Blanco. Cardiac output distribution in response to hypoxia in the chick embryo in the second half of the incubation time. *JOURNAL OF PHYSIOLOGY-LONDON*. 508:281-287, 1998.
- Mulder, T. L. M., J. C. vanGolde, F. W. Prinzen and C. E. Blanco. Cardiac output distribution in the chick embryo from stage 36 to 45. *Cardiovascular Research*. 34:525-528, 1997.
- MunchEllingsen, J., E. Bugge and K. Ytrehus. Blockade of the K-ATP-channel by glibenclamide aggravates ischemic injury, and counteracts ischemic preconditioning. *Basic Research In Cardiology*. 91:382-388, 1996.
- Murata, Y., K. Fujimori, E. J. Quilligan, N. Nagata, S. Ibara, T. Hirano and T. Kamimura. Cardiac oxygenation by extracorporeal membrane oxygenation in exteriorized fetal lambs. *American Journal Of Obstetrics And Gynecology*. 174:864-870, 1996.
- Nabavi, D. G., A. Cenic, J. Dool, R. M. L. Smith, F. Espinosa, R. A. Craen, A. W. Gelb and T. Y. Lee. Quantitative assessment of cerebral hemodynamics using CT: Stability, accuracy, and precision studies in dogs. *JOURNAL-OF-COMPUTER-ASSISTED-TOMOGRAPHY*. 234:506-515, 1999.
- Nagai, H., T. Nakashima, T. Suzuki and N. Yanagita. Effect of increased middle ear pressure on blood flow to the middle ear, inner ear and facial nerve in guinea pigs. *Acta Oto Laryngologica*. 116:439-442, 1996.
- Nakai, S., K. Ishikawa, I. Ogawa, H. Koka, N. Kamata, H. Akiyama and R. Katori. New collateral flow increasing early after coronary occlusion prevented myocardial necrosis in dogs. *Heart Vessels*. 10:171-7, 1995.
- Nakashima, T., K. Asami, K. Akanabe, K. Kuno, H. Nagai and N. Yanagita. Effect of carbon dioxide on facial nerve blood flow in rabbits. *Acta Otolaryngol Stockh*. 115:40-3., 1995.
- Nakashima, T., T. Suzuki, H. Morisaki and N. Yanagita. Blood flow in the cochlea, vestibular apparatus and facial nerve. *Acta Otolaryngol Stockh*. 111:738-42, 1991.

Nakashima, T., T. Suzuki and N. Yanagita. Cochlear blood flow under increased inner ear pressure. *Ann Otol Rhinol Laryngol.* 100:394-7., 1991.

Naredi, P., J. Mattson, L. Hafstrom and L. Jacobsson. Evaluation of blood flow measurements with microspheres and rubidium--an experimental study in rats. *Int J Microcirc Clin Exp.* 9:423-37, 1990.

Nellis, S. H., K. L. Carroll and A. M. Eggleston. Measurement of phasic velocities in vessels of intact freely beating hearts. *Am J Physiol.* 260:H1264-75, 1991.

Neumann, T. and G. Heusch. Myocardial, skeletal muscle, and renal blood flow during exercise in conscious dogs with heart failure. *American Journal Of Physiology Heart And Circulatory Physiology.* 42:H2452-H2457, 1997.

Nohara, R., K. Okuda, M. Ogino, R. Hosokawa, N. Tamaki, J. Konishi, Y. Fujibayashi, Y. Yonekura, M. Fujita and S. Sasayama. Evaluation of myocardial viability with iodine-123-BMIPP in a canine model. *Journal Of Nuclear Medicine.* 37:1403-1407, 1996.

Nose, Y., T. Nakamura and M. Nakamura. The microsphere method facilitates statistical assessment of regional blood flow. *Basic Res Cardiol.* 80:417-29, 1985.

Okuda, K., R. Nohara, M. Fujita, N. Tamaki, J. Konishi and S. Sasayama. Technetium-99m-pyrophosphate uptake as an indicator of myocardial injury without infarct. *J Nucl Med.* 35:1366-70., 1994.

Oohara, K., A. Usui, M. Murase, M. Tanaka and T. Abe. Regional cerebral tissue blood flow measured by the colored microsphere method during retrograde cerebral perfusion. *J Thorac Cardiovasc Surg.* 109:772-9., 1995.

Oohara, K., A. Usui, M. Tanaka, T. Abe and M. Murase. Determination of organ blood flows during retrograde inferior vena caval perfusion. *Ann Thorac Surg.* 58:139-45., 1994.

Ooiwa, H., T. Miura, T. Iwamoto, T. Ogawa, R. Ishimoto, T. Adachi and O. Iimura. Superoxide dismutase attenuated post-ischaemic contractile dysfunction in a myocardial xanthine oxidase deficient species. *Clin Exp Pharmacol Physiol.* 19:119-25., 1992.

Orgul, S., G. A. Cioffi, D. R. Bacon, A. Bhandari and E. M. VanBuskirk. Measurement of optic nerve blood flow with nonradioactive colored microspheres in rabbits. *Microvascular Research.* 51:175-186, 1996.

Orgul, S., G. A. Cioffi, D. R. Bacon and E. M. VanBuskirk. An endothelin-1-induced model of chronic optic nerve ischemia in rhesus monkeys. *Journal Of Glaucoma.* 5:135-138, 1996.

Orgul, S., G. A. Cioffi, D. J. Wilson, D. R. Bacon and E. M. VanBuskirk. An endothelin-1 induced model of optic nerve ischemia in the rabbit. *Investigative Ophthalmology & Visual Science.* 37:1860-1869, 1996.

Pinkerton, K. E., J. T. Gallen, R. R. Mercer, V. C. Wong, C. G. Plopper and B. K. Tarkington. Aerosolized fluorescent microspheres detected in the lung using confocal scanning laser microscopy. *Microsc Res Tech.* 26:437-43, 1993.

Pisarri, T. E., H. M. Coleridge and J. C. Coleridge. Reflex bronchial vasodilation in dogs evoked by injection of a small volume of water into a bronchus. *J Appl Physiol.* 75:2195-202., 1993.

Polissar, N. L., D. Stanford and R. W. Glenny. The 400 microsphere per piece "rule" does not apply to all blood flow studies. *Am. J. Physiol.* in press.

Powers, K. M., C. Schimmel, R. W. Glenny and C. M. Bernards. Cerebral blood flow determinations using fluorescent microspheres: variations on the sedimentation method validated. *Journal of Neuroscience Methods.* 87:159-165, 1999.

Preckel, B., W. Schlack, D. Obal, H. Barthel, D. Ebel, S. Grunert and V. Thamer. Effect of acidotic blood reperfusion on reperfusion injury after coronary artery occlusion in the dog heart. *JOURNAL OF CARDIOVASCULAR PHARMACOLOGY.* 31:179-186, 1998.

Presson, R. G., Jr., J. A. Graham, C. C. Hanger, P. S. Godbey, S. A. Gebb, R. A. Sidner, R. W. Glenny and W. W. Wagner, Jr. Distribution of pulmonary capillary red blood cell transit times. *J Appl Physiol.* 79:382-8, 1995.

Pries, A. R., K. Ley, M. Claassen and P. Gaehtgens. Red cell distribution at microvascular bifurcations. *Microvasc Res.* 38:81-101, 1989.

Prinzen, F. W. and R. W. Glenny. Developments in non-radioactive microsphere techniques for blood flow measurement. *Cardiovasc Res.* 28:1467-75, 1994.

Quinones, A. and B. J. Cheirif. New perspectives for perfusion imaging in echocardiography. *Circulation.* 83:III104-10, 1991.

Raab, S., E. Thein, A. G. Harris and K. Messmer. A new sample-processing unit for the fluorescent microsphere method. *American Journal of Physiology-Heart and Circulatory Physiology.* 45:H1801-H1806, 1999.

Rao, P. V., R. F. Stahl, B. R. Soller, K. G. Shortt, C. Hsi, K. J. Cotter, J. M. Belle Isle and J. M. Moran. Retrograde abdominal visceral perfusion: Is it beneficial? *Annals Of Thoracic Surgery.* 60:1704-1708, 1995.

Rasmussen, O., F. F. Lauszus, C. Christiansen, C. Thomsen and K. Hermansen. Differential Effects Of Saturated and Monounsaturated Fat On Blood Glucose and Insulin Responses In Subjects With Non-Insulin-Dependent Diabetes Mellitus. *American Journal of Clinical Nutrition.* 63:249-253, 1996.

Rattigan, S., G. J. Appleby, K. A. Miller, J. T. Steen, K. A. Dora, E. Q. Colquhoun and M. G. Clark. Serotonin inhibition of 1-methylxanthine metabolism parallels its vasoconstrictor activity and inhibition of oxygen in perfused rat hindlimb. *Acta Physiologica Scandinavica.* 161:161-169, 1997.

Reeves, W. J. and K. Rakusan. Myocardial capillary flow pattern as determined by the method of coloured microspheres. *Adv. Exp. Med. Biol.* 222:447-53, 1988.

Revelly, J. P., T. Ayuse, N. Brienza, H. E. Fessler and J. L. Robotham. Endotoxic shock alters distribution of blood flow within the intestinal wall. *Critical Care Medicine*. 24:1345-1351, 1996.

Ring, G. C., A. S. Blum, T. Kurbatov, W. G. Moss and W. Smith. Size of microspheres passing through pulmonary circuit in the dog. *American Journal of Applied Physiology*. 200:1191-1196, 1961.

Robertson, H. T., R. W. Glenny, D. Stanford, L. M. McInnes, D. L. Luchtel and D. Covert. High-resolution maps of regional ventilation utilizing inhaled fluorescent microspheres. *Journal Of Applied Physiology*. 82:943-953, 1997.

Rolf, N., A. Meissner, H. VanAken, T. P. Weber, D. Hammel and T. Mollhoff. The effects of thoracic epidural anesthesia on functional recovery from myocardial stunning in propofol-anesthetized dogs. *Anesthesia And Analgesia*. 84:723-729, 1997.

Rolf, N., M. VandeVelde, P. F. Wouters, T. Mollhoff, T. P. Weber and H. K. VanAken. Thoracic epidural anesthesia improves functional recovery from myocardial stunning in conscious dogs. *Anesthesia And Analgesia*. 83:935-940, 1996.

Rose, J., T. Ehring, S. G. Sakka, A. Skyschally and G. Heusch. Aspirin does not prevent the attenuation of myocardial stunning by the ACE inhibitor ramipril. *Journal Of Molecular And Cellular Cardiology*. 28:603-613, 1996.

Rovainen, C. M., T. A. Woolsey, N. C. Blocher, D. B. Wang and O. F. Robinson. Blood flow in single surface arterioles and venules on the mouse somatosensory cortex measured with videomicroscopy, fluorescent dextrans, nonoccluding fluorescent beads, and computer-assisted image analysis. *J Cereb Blood Flow Metab*. 13:359-71, 1993.

Rudis, E., R. N. Gates, H. Laks, D. C. Drinkwater, A. Ardehali, A. Aharon and P. Chang. Coronary sinus ostial occlusion during retrograde delivery of cardioplegic solution significantly improves cardioplegic distribution and efficacy [see comments]. *J Thorac Cardiovasc Surg*. 109:941-6; discussion 946-7., 1995.

Rudolph, A. M. and M. A. Heymann. The circulation of the fetus in utero. Methods for studying distribution of blood flow, cardiac output and organ blood flow. *Circ Res*. 21:163-84, 1967.

Saida, A., H. Ito, T. Shibuya and Y. Watanabe. Time-course alterations of monoamine levels and cerebral blood flow in brain regions after subarachnoid hemorrhage in rats. *Brain Research Bulletin*. 43:69-80, 1997.

Saitou, H., T. Watanabe, J. W. Zhang, N. Oshikiri, Y. Iijima, K. Inui, S. Kuraoka and Y. Shimazaki. Regional tissue blood flow and pH in the brain during deep hypothermic retrograde brain perfusion. *Journal Of Surgical Research*. 72:135-140, 1997.

Sakaki, M., Y. Taenaka, E. Tatsumi, T. Nakatani, M. Kinoshita, H. Akagi, T. Masuzawa, Y. Matsuo, K. Inoue, Y. Baba and et al. Pulmonary function in a non-pulsatile pulmonary circulation. *Asao J.* 38:M366-9., 1992.

Sakka, S. G., D. R. Wallbridge and G. Heusch. Methods for the measurement of coronary blood flow and myocardial perfusion - Glossary. *Basic Research In Cardiology*. 91:155-178, 1996.

Sakurai, H., L. D. Traber and D. L. Traber. Altered systemic organ blood flow after combined injury with burn and smoke inhalation. *SHOCK*. 9:369-374, 1998.

Salmons, S., A. T. M. Tang, J. C. Jarvis, H. Degens, M. Hastings and T. L. Hooper. Morphological and functional evidence, and clinical importance, of vascular anastomoses in the latissimus dorsi muscle of the sheep. *JOURNAL OF ANATOMY*. 193:93-104, 1998.

Sato, M., T. Saito, M. Mitsugi, S. I. Saitoh, T. Niitsuma, K. Maehara and Y. Maruyama. Effects of cardiac contraction and coronary sinus pressure elevation on collateral circulation. *American Journal Of Physiology Heart And Circulatory Physiology*. 40:H1433-H1440, 1996.

Saxena, P. R. and P. D. Verdouw. Tissue blood flow and localization of arteriovenous anastomoses in pigs with microspheres of four different sizes. *Pflugers Arch.* 403:128-35, 1985.

Schenarts, P. J., H. G. Bone, L. D. Traber and D. L. Traber. Effect of severe smoke inhalation injury on systemic microvascular blood flow in sheep. *Shock*. 6:201-205, 1996.

Schimmel, C., D. Frazer and R.W. Glenny. Extending Flourescent Microsphere Methods for Regional Organ Blood Flow to 13 Simultaneous Colors. *Am J Physiol Heart Circ Physiol*. 280: H2496-506, 2001.

Schimmel, C., D. Frazer, S.R. Huckins and R.W. Glenny. Validation of Automated Spectrofluorimetry for Measurement of Regional Organ Perfusion Using Fluorescent Microspheres. *Comput Methods Programs Biomed*. 62:115-25, 2000.

Schlack, W., D. Ebel, S. Grunert, S. Halilovic, O. Meyer and V. Thamer. Effect of heart rate reduction by 4-(N-ethyl-N-phenylamino)-1,2-dimethyl-6- (methylamino)pyrimidinium chloride on infarct size in dog. *ARZNEIMITTEL-FORSCHUNG/DRUG RESEARCH*. 48:26-33, 1998.

Schmid-Schonbein, G. W., R. Skalak, S. Usami and S. Chien. Cell distribution in capillary networks. *Microvasc Res*. 19:18-44, 1980.

Schosser, R., K. E. Arfors and K. Messmer. MIC-II - a program for the determination of cardiac output, arterio-venous shunt and regional blood flow using the radioactive microsphere method. *Comput Programs Biomed*. 9:19-38, 1979.

Schwanke, U., H. Strauss, G. Arnold and J. D. Schipke. Analysis of respiratory water - A new method for evaluation of myocardial energy metabolism. *J Appl Physiol*. 81:2115-2122, 1996.

Shabsigh, A., D. T. Chang, D. F. Heitjan, A. Kiss, C. A. Olsson, P. J. Puchner and R. Buttyan. Rapid reduction in blood flow to the rat ventral prostate gland after castration: Preliminary evidence that androgens influence prostate size by regulating blood flow to the prostate gland and prostatic endothelial cell survival. *PROSTATE*. 36:201-206, 1998.

Shimada, K., K. Yoshida, H. Tadokoro, S. Kitsukawa, A. Takami, K. Suzuki, S. Tanada and Y. Masuda. High-resolution cardiac PET in rabbits: Imaging and quantitation of myocardial blood flow. *Journal of Nuclear Medicine*. 39:2022-2027, 1998.

Shin, C., M. P. Kinsky, J. A. Thomas, D. L. Traber and G. C. Kramer. Effect of cutaneous burn injury and resuscitation on the cerebral circulation in an ovine model. *BURNS*. 24:39-45, 1998.

Sidi, A. and W. Rush. An alternative to radioactive microspheres for measuring regional myocardial blood flow .1. colored microspheres. *Journal of Cardiothoracic & Vascular Anesthesia*. 10:368-373, 1996.

Sidi, A. and W. Rush. An alternative to radioactive microspheres for measuring regional myocardial blood flow .2. laser-doppler perfusion monitor. *Journal of Cardiothoracic & Vascular Anesthesia*. 10:374-377, 1996.

Simor, T., W. J. Chu, L. Johnson, A. Safranko, M. Doyle, G. M. Pohost and G. A. Elgavish. In vivo MRI visualization of acute myocardial ischemia and reperfusion in ferrets by the persistent action of the contrast agent Gd(BME-DTTA). *Circulation*. 92:3549-3559, 1995.

Smith, M. V., R. F. Stahl, C. Cronin, C. Hsi, J. M. Li, J. BelleIsle, M. Knox and T. J. Vander-Salm. Retrograde coronary sinus perfusion provides non-homogeneous myocardial blood flow. *Cardiology*. 83:331-8, 1993.

Souders, J. E., S. C. George, N. L. Polissar, E. R. Swenson and M. P. Hlastala. Tracheal gas exchange: perfusion-related differences in inert gas elimination. *J Appl Physiol*. 79:918-28, 1995.

Spinale, F. G., R. Tanaka, F. A. Crawford and M. R. Zile. Changes in myocardial blood flow during development of and recovery from tachycardia-induced cardiomyopathy. *Circulation*. 85:717-29, 1992.

Steinhausen, M., B. Zimmerhackl, H. Thederan, R. Dussel, N. Parekh, H. U. Esslinger, G. von-Hagens, D. Komitowski and F. D. Dallenbach. Intraglomerular microcirculation: measurements of single glomerular loop flow in rats. *Kidney Int*. 20:230-9, 1981.

Stryer, L. and R. P. Haugland. Energy transfer: A spectroscopic ruler. *Proc. Natnl. Acad. Sci.* 58:719-726, 1967.

Suckfull, M., S. Holtmann and R. Hecht. A new method for measuring internal ear circulation using colored microspheres. *Laryngorhinootologie*. 74:141-4., 1995.

Suckfull-M., Winkler-G., Thein-E., Raab-S., Schorn-K. and Mees-K. Changes in serum osmolarity influence the function of outer hair cells. *Acta Oto-Laryngologica*. 119:316-321, 1999.

Suo, Z. M., J. Humphrey, A. Kundtz, F. Sethi, A. Placzek, F. Crawford and M. Mullan. Soluble Alzheimers beta-amyloid constricts the cerebral vasculature in vivo. *Neuroscience Letters*. 257:77-80, 1998.

Suzuki, T., T. Nakashima and N. Yanagita. Effects of increased cerebrospinal fluid pressure on cochlear and cerebral blood flow. *Eur Arch Otorhinolaryngol*. 250:332-6, 1993.

Svensson, A. M., C. G. Ostenson, S. Sandler, S. Efendic and L. Jansson. Inhibition of nitric oxide synthase by NG-nitro-L-arginine causes a preferential decrease in pancreatic islet blood flow in normal rats and spontaneously diabetic GK rats. *Endocrinology*. 135:849-53., 1994.

Symons, J. D., S. D. Correa and S. Schaefer. Na-H exchange inhibition with cariporide limits functional impairment caused by repetitive ischemia. *Journal of Cardiovascular Pharmacology*. 32:853-862, 1998.

Tabor, O. B., M. J. Bosse, K. G. Greene, H. E. Gruber, K. Kaysinger, S. H. Sims, S. R. Blumenthal and J. E. Kellam. Effects of surgical approaches for acetabular fractures with associated gluteal vascular injury. *JOURNAL OF ORTHOPAEDIC TRAUMA*. 12:78-84, 1998.

Takahata, O., K. Ichihara, Y. Abiko and H. Ogawa. Sevoflurane preserves endocardial blood flow during coronary ligation in dogs: comparison with adenosine. *ACTA ANAESTHESIOLOGICA SCANDINAVICA*. 42:225-231, 1998.

Takanashi, H. and M. Akima. Protective effects of arbabiprostil against indomethacin-induced gastric lesions in rats: significance of maintained gastric blood flow. *Jpn J Pharmacol*. 59:349-55., 1992.

Takao, S., M. Maeda, M. Inoue, T. Fukushima, M. Tomonaga and Y. Hayashida. Spinal cord blood flow decreases following microinjection of sodium nitroprusside into the nucleus tractus solitarii of anesthetized rats. *Neuroscience Research*. 25:285-291, 1996.

Tamaki, Y., E. Kawamoto, M. Araie, S. Eguchi and H. Fujii. [Noninvasive two-dimensional analysis of retinal microcirculation using the laser speckle phenomenon--II. Comparison of results with those obtained with the microsphere technique]. *Nippon Ganka Gakkai Zasshi*. 98:169-74., 1994.

Tan, D. Y., M. R. Cernadas, P. Aragoncillo, M. A. Castilla, M. V. A. Arroyo, A. J. L. Farre, S. Casado and C. Caramelo. Role of nitric oxide-related mechanisms in renal function in ageing rats. *NEPHROLOGY DIALYSIS TRANSPLANTATION*. 13:594-601, 1998.

Tan, W. P., K. W. Riggs, R. L. Thies and D. W. Rurak. Use of an automated fluorescent microsphere method to measure regional blood flow in the fetal lamb. *Canadian Journal Of Physiology And Pharmacology*. 75:959-968, 1997.

Tanabe, T., T. Iwamoto, O. Iwata, M. Aikawa, S. Kusuzaki, S. Handa, Y. Shinozaki and H. Mori. Electrophysiologic and blood-flow responses in the endocardium and epicardium to disopyramide and MS-551 during myocardial ischemia in the dog. *JOURNAL-OF-CARDIOVASCULAR-PHARMACOLOGY*. 34:275-286, 1999.

Tang, A. T. M., J. C. Jarvis, T. L. Hooper and S. Salmons. Observation and basis of improved blood flow to the distal latissimus dorsi muscle: a case for electrical stimulation prior to grafting. *Cardiovascular Research*. 40:131-137, 1998.

Tatsumi, E., K. Miyazaki, K. Toda, Y. Taenaka, T. Nakatani, Y. Baba, T. Masuzawa, Y. Wakisaka, K. Eya, T. Nishimura, Y. Takewa, T. Ohno and H. Takano. Influence of non pulsatile systemic circulation on tissue blood flow and oxygen metabolism. *Asao Journal*. 42:M757-M762, 1996.

Tokuda, N. and R. B. Levy. 1,25-dihydroxyvitamin D-3 stimulates phagocytosis but suppresses HLA-DR and CD13 antigen expression in human mononuclear phagocytes. *Proceedings Of The Society For Experimental Biology And Medicine*. 211:244-250, 1996.

Tretli, S. and M. Gaard. Lifestyle Changes During Adolescence and Risk Of Breast Cancer - an Ecologic Study Of the Effect Of World War II In Norway. *Cancer Causes & Control*. 7:507-512, 1996.

Tsuchida, Y., N. Aoki, O. Fukuda, M. Nakano and H. Igarashi. Changes in hemodynamics in jejunal flaps of rabbits due to ischemia, venous congestion, and reperfusion as measured by means of colored microspheres., 1998

Tsumoto, S., S. Kawashima and T. Iwasaki. Unaltered size of right ventricular infarct in dogs with right ventricular hypertrophy induced by pressure overload. *Am J Physiol*. 268:H1781-7, 1995.

Usui, K., T. Tanabe, S. Handa, Y. Shinozaki and H. Mori. Disproportional response between refractory period and blood flow to alpha(1) and beta-andrenoceptor blockade in canine ischemic myocardium. *Cardiovascular Drugs and Therapy*. 12:561-571, 1998.

Utley, J., E. L. Carlson, J. I. Hoffman, H. M. Martinez and G. D. Buckberg. Total and regional myocardial blood flow measurements with 25 micron, 15 micron, 9 micron, and filtered 1-10 micron diameter microspheres and antipyrine in dogs and sheep. *Circ Res*. 34:391-405, 1974.

van Golde, J. M., T. A. Mulder, E. Scheve, F. W. Prinzen and C. E. Blanco. Hyperoxia and local organ blood flow in the developing chick embryo. *J Physiol (Lond)*. 515:243-248, 1999.

van Oosterhout, M. F., H. M. Willigers, R. S. Reneman and F. W. Prinzen. Fluorescent microspheres to measure organ perfusion: validation of a simplified sample processing technique. *Am J Physiol*. 269:H725-33, 1995.

van Oosterhout, M. F. M. and F. W. Prinzen. The fluorescent microsphere method for determination of organ blood flow. *FASEB J*. 8:A854, 1994.

van Oosterhout, M. F. M., F. W. Prinzen, S. Sakurada, R. W. Glenny and J. R. S. Hales. Fluorescent microspheres are superior to radioactive microspheres in chronic blood flow measurements. *AMERICAN JOURNAL OF PHYSIOLOGY-HEART AND CIRCULATORY PHYSIOLOGY*. 44:H110-H115, 1998.

Vandevska-Radunovic, V., A. B. Kristiansen, K. J. Heyeraas and S. Kvinnslund. Changes in blood circulation in teeth and supporting tissues incident to experimental tooth movement. *Eur J Orthod*. 16:361-9, 1994.

Vandevska-Radunovic, V., I. H. Kvinnslund and S. Kvinnslund. Effect of inferior alveolar nerve axotomy on periodontal and pulpal blood flow subsequent to experimental tooth movement in rats. *ACTA ODONTOLOGICA SCANDINAVICA*. 56:57-64, 1998.

vanDoorn, C. A. M., H. Degens, M. S. Bhabra, C. B. W. Till, T. E. Shaw, J. C. Jarvis, S. Salmons and T. L. Hooper. Intramural blood flow of skeletal muscle ventricles functioning as aortic counterpulsators. *Annals Of Thoracic Surgery*. 64:86-93, 1997.

Vescovo, G., C. Ceconi, P. Bernocchi, R. Ferrari, U. Carraro, G. B. Ambrosio and L. DallaLibera. Skeletal muscle myosin heavy chain expression in rats with monocrotaline-induced cardiac hypertrophy and failure. Relation to blood flow and degree of muscle atrophy. *CARDIOVASCULAR RESEARCH*. 39:233-241, 1998.

Viroonchatapan, E., H. Sato, M. Ueno, I. Adachi, K. Tazawa and I. Horikoshi. Magnetic targeting of thermosensitive magnetoliposomes to mouse livers in an in situ on-line perfusion system. *Life Sciences*. 58:2251-2261, 1996.

Walker, D. M., J. M. Walker and D. M. Yellon. Global myocardial ischemia protects the myocardium from subsequent regional ischemia. *Cardioscience*. 4:263-6., 1993.

Walland, A., H. Weihs and E. Mutschler. Perfusion pressure and transmural flow distribution in the left ventricles of isolated rat hearts. *Clin Exp Pharmacol Physiol*. 20:723-9, 1993.

Walter, B., R. Bauer, E. Gaser and U. Zwiener. Validation of the multiple colored microsphere technique for regional blood flow measurements in newborn piglets. *Basic Research In Cardiology*. 92:191-200, 1997.

Walter, D. H., U. Hink, T. Asahara, E. VanBelle, J. Horowitz, Y. Tsurumi, R. Vandlen, H. Heinsohn, B. Keyt, N. Ferrara, J. F. Symes and J. M. Isner. The in vivo bioactivity of vascular endothelial growth factor vascular permeability factor is independent of N-linked glycosylation. *Laboratory Investigation*. 74:546-556, 1996.

Walther, S., K. Domino, R. Glenny, N. Polissar and M. Hlastala. Pulmonary blood flow distribution has a hilar-to-peripheral gradient in awake, prone sheep. *J. Appl. Physiol.* 82:678-685, 1997.

Walther, S. M., K. B. Domino, R. W. Glenny and M. P. Hlastala. Pulmonary blood flow distribution in sheep: Effects of anesthesia, mechanical ventilation, and change in posture. *Anesthesiology*. 87:335-342, 1997.

Walther, S. M., K. B. Domino, R. W. Glenny and M. P. Hlastala. Positive end-expiratory pressure redistributes perfusion to dependent lung regions in supine but not in prone lambs. *Critical Care Medicine*. 27:37-45, 1999.

Wang, X. J., F. Q. Li, S. Said, J. M. Capasso and A. M. Gerdes. Measurement of regional myocardial blood flow in rats by unlabeled microspheres and Coulter Channelyzer. *American Journal Of Physiology Heart And Circulatory Physiology*. 40:H1656-H1665, 1996.

Weihs, H., E. Mutschler and A. Walland. Alinidine prevents hypoperfusion-induced transmural flow redistribution in isolated paced rat hearts. *Clin Exp Pharmacol Physiol*. 21:471-6, 1994.

Wiedeman, M. P. Dimensions of Blood Vessels from Distributing Artery to Collecting Vein. *Circulation Research*. XII:375-378, 1963.

Wieland, W., P. F. Wouters, H. Van Aken and W. Flameng. Measurement of Organ Blood Flow with Coloured Microspheres: a First Time-Saving Improvement using Automated Spectrophotometry. *Proceedings of Computers in Cardiology*. 691-694, 1993.

Wouters, P. F., M. VandeVelde, H. VanAken and W. Flameng. Ischemic event characteristics determine the extent of myocardial stunning in conscious dogs. *Basic Research In Cardiology*. 91:140-146, 1996.

Wu, C. H., D. C. Lindsey, D. L. Traber, C. E. Cross, D. N. Herndon and G. C. Kramer. Measurement of bronchial blood flow with radioactive microspheres in awake sheep. *J Appl Physiol*. 65:1131-9, 1988.

Xuan, B., Y. H. Zhou, R. Varma and G. C. Y. Chiou. Effects of some N-nitropyrazole derivatives on ocular blood flow and retinal function recovery after ischemic insult. *Journal of Ocular Pharmacology and Therapeutics*. 15:135-142, 1999.

Yamamoto, N., T. Kohmoto, A. G. Gu, C. DeRosa, C. R. Smith and D. Burkhoff. Angiogenesis is enhanced in ischemic canine myocardium by transmyocardial laser revascularization. *JOURNAL OF THE AMERICAN COLLEGE OF CARDIOLOGY*. 31:1426-1433, 1998.

Yamasaki, T. and T. Tsutsui. Spontaneous emission from fluorescent molecules embedded in photonic crystals consisting of polystyrene microspheres. *APPLIED PHYSICS LETTERS*. 72:1957-1959, 1998.

Yanagi, S., K. Takeuchi, T. Takeda, M. Ishikawa and I. Miura. Effects of inotropic stimulation on phosphate compounds in ischaemic canine hearts. *Cardiovasc Res*. 27:1435-43., 1993.

Yanagi, S., K. Takeuchi, T. Takeda, M. Ishikawa and I. Miura. Comparison of the effects of dobutamine and isoproterenol in ischemic hearts by phosphorus-31 nuclear magnetic resonance spectroscopy. *Jpn Circ J*. 58:338-50., 1994.

Yang, R., Q. Liu, J. L. Unthank, M. D. Pescovitz and J. L. Grosfeld. Preservation of postprandial intestinal hyperemic response after small bowel transplantation. *J Pediatr Surg*. 30:1090-2., 1995.

Yen, R. T. and Y. C. Fung. Effect of velocity of distribution on red cell distribution in capillary blood vessels. *J R Coll Gen Pract Occas Pap*. 235:H251-7, 1978.

Yipintsoi, T., W. A. Dobbs, Jr., P. D. Scanlon, T. J. Knopp and J. B. Bassingthwaigte. Regional distribution of diffusible tracers and carbonized microspheres in the left ventricle of isolated dog hearts. *Circ Res*. 33:573-87, 1973.

Zeltner, T. B., T. D. Sweeney, W. A. Skornik, H. A. Feldman and J. D. Brain. Retention and clearance of 0.9-micron particles inhaled by hamsters during rest or exercise. *J Appl Physiol.* 70:1137-45, 1991.

Zink, B. J., C. H. Schultz, X. Wang, M. Mertz, S. A. Stern and A. L. Betz. Effects of ethanol on brain lactate in experimental traumatic brain injury with hemorrhagic shock. *BRAIN-RESEARCH.* 837:1-7, 1999.

Zippel, K. C., H. B. Lillywhite and C. R. J. Mladinich. Contribution of the vertebral artery to cerebral circulation in the rat snake *Elaphe obsoleta*. *JOURNAL OF MORPHOLOGY.* 238:39-51, 1998.

Zwissler, B., R. Schosser, C. Weiss, V. Iber, M. Weiss, C. Schwickert, P. Spengler and K. Messmer. Methodological error and spatial variability of organ blood flow measurements using radiolabeled microspheres. *Res Exp Med Berl.* 191:47-63, 1991.

